



## PPD Administration Record

**PATIENT NAME:** \_\_\_\_\_ **DATE:** \_\_\_\_\_

**FACILITY:** \_\_\_\_\_

**FACILITY ADDRESS:** \_\_\_\_\_

1. **Initial Test Dose:** Tuberculin Skin Test 0.1 ml (5TU) PPD Mantoux

Date given: \_\_\_\_\_ Lot #: \_\_\_\_\_ Expiration Date: \_\_\_\_\_

Time given: \_\_\_\_\_ Site: Rt. Forearm: \_\_\_\_\_ Lt. Forearm: \_\_\_\_\_

Administered by: \_\_\_\_\_

Date read: \_\_\_\_\_ Time: \_\_\_\_\_ Inuduration: \_\_\_\_\_

Erythema: \_\_\_\_\_ mm Edema: \_\_\_\_\_ mm Necrosis: \_\_\_\_\_ mm

\_\_\_\_\_  
Name of person reading test (printed)

\_\_\_\_\_  
Signature of person reading test

\_\_\_\_\_  
Position held of person reading the test

\_\_\_\_\_  
Date

2. **Second Test Dose** (*give only if 2 step test ordered*)

**Second Test Dose:** 5TU PPD Mantoux if initial test dose negative

Give 7-14 days after initial test dose if 2 step test ordered

Date given: \_\_\_\_\_ Lot #: \_\_\_\_\_ Expiration Date: \_\_\_\_\_

Time given: \_\_\_\_\_ Site: Rt. Forearm: \_\_\_\_\_ Lt. Forearm: \_\_\_\_\_

Administered by: \_\_\_\_\_

Date read: \_\_\_\_\_ Time: \_\_\_\_\_ Inuduration: \_\_\_\_\_

Erythema: \_\_\_\_\_ mm Edema: \_\_\_\_\_ mm Necrosis: \_\_\_\_\_ mm

\_\_\_\_\_  
Name of person reading test (printed)

\_\_\_\_\_  
Signature of person reading test

\_\_\_\_\_  
Position held of person reading the test

\_\_\_\_\_  
Date

**Interpretation of test:** 48-72 hours after administration

**Positive:** palpable induration of 10mm or more 5mm if had close contact with individuals with tuberculosis

**Doubtful Reaction:** 5-9 mm, retest second site with 5TU

**Negative Reaction:** induration of less than 5mm to both first and second test